

Conflict-Free Coloring for Rectangle Ranges Using $\tilde{O}(n^{382+\epsilon})$ Colors

Deepak Ajwani * Khaled Elbassioni * Sathish Govindarajan *
Saurabh Ray †

Abstract

Given a set of points $P \subseteq \mathbb{R}^2$, a *conflict-free coloring* of P is an assignment of colors to points of P , such that each non-empty axis-parallel rectangle T in the plane contains a point whose color is distinct from all other points in $P \cap T$. This notion has been the subject of recent interest, and is motivated by frequency assignment in wireless cellular networks: one naturally would like to minimize the number of frequencies (colors) assigned to bases stations (points), such that within any range (for instance, rectangle), there is no interference. We show that any set of n points in \mathbb{R}^2 can be conflict-free colored with $\tilde{O}(n^{382+\epsilon})$ colors in expected polynomial time, for any arbitrarily small $\epsilon > 0$. This improves upon the previously known bound of $O(\sqrt{n \log \log n / \log n})$.

Keywords: Computational geometry, Combinatorial geometry, Conflict-free coloring

*Max Planck Institut für Informatik, Saarbrücken, Germany. E-mail: {ajwani, elbassio, sgovinda}@mpi-sb.mpg.de

†Dept. of Computer Science, Universität des Saarlandes, Saarbrücken, Germany. E-mail: saurabh@cs.uni-sb.de

1 Introduction

The study of conflict-free coloring is motivated by the frequency assignment problem in wireless networks. A wireless network is a heterogeneous network consisting of *base stations* and *agents*. The base stations have a fixed location, and are interlinked via a fixed backbone network, while the agents are typically mobile and can connect to the base stations via radio links. The base stations are assigned fixed frequencies to enable links to agents. The agents can connect to any base station, provided that the radio link to that particular station has good reception. Good reception is only possible if i) the base station is located within range, and ii) no other base station within range of the agent has the same frequency assignment (to avoid interference). Thus the fundamental problem of frequency-assignment in cellular networks is to assign frequencies to base stations, such that an agent can always find a base station with unique frequency among the base stations in its range. Naturally, due to cost, flexibility and other restrictions, one would like to minimize the total number of assigned frequencies.

The study of the above problem was initiated in [ELRS03], and continued in a series of recent papers [PT03, HS05, FLM⁺05, AS06, BNCS06, CKS06, Che06, EM06, Smo06]. It can be formally described as follows. Let $P \subseteq \mathbb{R}^2$ be a set of points and \mathcal{R} be a set of ranges (e.g. the set of all discs or rectangles in the plane). A *conflict-free* coloring (CF-coloring in short) of P w.r.t. the range \mathcal{R} is an assignment of a color to each point $p \in P$ such that for any range $T \in \mathcal{R}$ with $T \cap P \neq \emptyset$, the set $T \cap P$ contains a point of unique color. Naturally, the goal is to assign a conflict-free coloring to the points of P with the *smallest* number of colors possible.

The work in [ELRS03] presented a general framework for computing a conflict-free coloring for several types of ranges. In particular, for the case where the ranges are discs in the plane, they present a polynomial time coloring algorithm that uses $O(\log n)$ colors for conflict-free coloring and this bound is shown to be tight. This result was then extended in [HS05] by considering the case where the ranges are axis-parallel rectangles in the plane. This seems much harder than the disc case, and the work in [HS05] presented a simple algorithm that uses $O(\sqrt{n})$ colors. As mentioned in [HS05], this can be further improved to $O(\sqrt{n \log \log n / \log n})$ using the sparse neighbourhood property of the conflict-free graph, as independently observed in [AKS99, PT03]. Currently, this is the best known upper bound for CF-coloring axis-parallel rectangles. A lower bound of $\Omega(\log n)$ trivially follows from the intervals case. Very recently, Pach and Toth showed that there exists a set of n points which need $\Omega(\log^2 n)$ colors for a conflict-free coloring.

Recent works have shown that one can obtain better upper bounds for special cases of this problem. In [HS05], it was shown that for the case of random points in a unit square, $O(\log^4 n)$ colors suffice, and for points lying in an *exact* uniform $\sqrt{n} \times \sqrt{n}$ grid, $O(\log n)$ colors are sufficient. Chen [Che06] showed that polylogarithmic number of colors suffice for the case of *nearly equal* rectangle ranges. Elbassioni and Mustafa [EM06] asked the following question: Given a set of points P in the plane, can we insert new points Q such that the conflict free coloring of $P \cup Q$ requires fewer colors? They showed that by inserting $O(n^{1-\epsilon})$ points, $P \cup Q$ can be conflict free colored using $\tilde{O}(n^{3(1+\epsilon)/8})$ colors.

While the CF-coloring problem is closed for disc ranges, the upper bounds are

very far from the currently known lower bounds for axis-parallel rectangular ranges. It remains a very interesting problem to reduce this gap between upper and lower bounds, and this is, in fact, the main open problem posed in [HS05]. In this paper, we improve the upper bound significantly.

Theorem 1.1 *Any set of n points in \mathbb{R}^2 can be conflict-free colored with respect to rectangle ranges using $\tilde{O}(n^{.382+\epsilon})$ colors, in expected polynomial time, for any arbitrarily small $\epsilon > 0$.*

Our main tool for proving this theorem is a probabilistic coloring technique, introduced in [EM06], that can be used to get a coloring with weaker properties, which we call *quasi-conflict-free* coloring. This will be combined with dominating sets, monotone sequences, and careful gridding of the pointset, in a recursive way, to obtain the claimed result. We start with some definitions and preliminaries in Section 2. To illustrate our ideas, we sketch a simple $\tilde{O}(n^{6/13})$ conflict free coloring algorithm in Section 3. We recall the quasi-conflict-free coloring technique in a slightly more general form in Section 4. The main algorithm will be given in Section 5 and analyzed in Section 6. Finally, we derive some corollaries of Theorem 1.1 in Section 7.

2 Preliminaries

By $\mathcal{R} \subseteq 2^{\mathbb{R}^2}$ we denote the set of all *axis-parallel* rectangles.

Definition 2.1 (Conflict-free coloring) *Let P be a set of points in \mathbb{R}^2 . A coloring of P is a function $\chi : P \mapsto \mathbb{N}^* \stackrel{\text{def}}{=} \mathbb{N} \cup \mathbb{N}^2 \cup \dots$ from P to the sequences of natural numbers. A rectangle $T \in \mathcal{R}$ is said to be conflict-free with respect to a coloring χ if either $T \cap P = \emptyset$, or there exists a point $p \in P \cap T$ such that $\chi(p) \neq \chi(p')$ for all $p \neq p' \in P \cap T$. A coloring χ is said to be conflict-free (w.r.t. \mathcal{R}) if every rectangle $T \in \mathcal{R}$ is conflict-free w.r.t. χ .*

Definition 2.2 (Dominating sets) *For a point $p = (p^x, p^y) \in \mathbb{R}^2$, define $W_1(p) = \{q \in \mathbb{R}^2 \mid q^x \geq p^x, q^y \geq p^y\}$ to be the upper right quadrant defined by p . Similarly, let $W_2(p)$, $W_3(p)$ and $W_4(p)$ be the upper left, lower right and lower left quadrants respectively. Define the dominating set of type i for a pointset $P \subseteq \mathbb{R}^2$, denoted by $D_i(P)$, $1 \leq i \leq 4$, as follows:*

$$D_i(P) = \{p \in P \mid W_i(p) \cap P = \emptyset\}$$

Definition 2.3 (Monotonic sets) *Let $P = \{p_1, p_2, \dots, p_k\}$ be a set of points that is sorted by x coordinate. P is monotonic non-decreasing (resp. monotonic non-increasing) if $p_j^y \geq p_i^y$ (resp. $p_j^y \leq p_i^y$) $\forall 1 \leq i, j \leq k, j > i$.*

It is easy to see that the dominating set of type 2 and 3 (resp. type 1 and 4) are monotonic non-decreasing (resp. non-increasing).

Definition 2.4 (r -Grid) *Let $P \subseteq \mathbb{R}^2$ be a set of points in the plane and $r \in \mathbb{Z}_+$ be a positive integer. An r -grid on P , denoted by $G_r = G_r(P)$, is an $r \times r$ axis-parallel*

grid containing all points of P . For $i = 1, \dots, r$, denote by R_i and C_i the subsets of P lying in the i th row and column of G_r , respectively. Denote respectively by $b(G_r)$ and $B(G_r)$, the minimum and maximum number of points of P in a row or a column of G_r . For $1 \leq h \leq 2r - 1$, let M_1^h (resp. M_h^2) be the set of grid cells lying along a diagonal h of positive slope (resp. negative slope) in G_r , and for $l = 1, \dots, 4$, let $\mathcal{D}_l^h = \cup_{(i,j) \in M_h} D_l(R_i \cap C_j)$ be the union of dominating sets of type l over grid cells in M_h .

Note that, for $l = 2, 3$ and $1 \leq h \leq 2r - 1$, \mathcal{D}_l^h is either monotonic non-decreasing, since the grid cells in M_h^1 , which lie along the diagonal of positive slope, are horizontally and vertically separated and hence the union of $D_l(R_i \cap C_j)$ (which are monotonic non-decreasing), is also monotonic non-decreasing. By similar argument, for $l = 1, 4$ with M_h^2 and $1 \leq h \leq 2r - 1$, \mathcal{D}_l^h is either monotonic non-increasing

When we talk about a grid $G_r(P)$ on a point set P , we may and will assume without loss of generality that no point of P lies on the vertical or horizontal lines defining the grid (otherwise a large monotone sequence exists, see Section 5 for more details). In particular, the grid lines define partitions of the pointset along the horizontal and vertical directions.

Definition 2.5 (Quasi-conflict-free coloring) *Given $P \subseteq \mathbb{R}^2$, and a grid $G_r = G_r(P)$ on P , we call a coloring $\chi : P \mapsto \mathbb{N}^*$ quasi-conflict-free with respect to G_r , if every axis-parallel rectangle which contains points only from the same row or the same column of G_r is conflict-free.*

Let G_r be a r -grid on a point set P such that $b(G_r) = B(G_r) = b$. It is shown in [EM06] that there exists a quasi-conflict-free coloring of $G_r(P)$ requiring $\tilde{O}(b^{3/4})$ colors.

3 An $\tilde{O}(n^{6/13})$ conflict free coloring algorithm

In this section, we sketch a simple algorithm for CF-coloring P in order to illustrate the main ideas. This algorithm CF-colors P using $\tilde{O}(n^{6/13})$ colors. We can assume w.l.o.g. that P has no monotone sequences of size $\Omega(n^{7/13})$. If there is one, we pick every other point in the sequence (this is a set I of size $\Omega(n^{7/13})$), color them all with one color, and recurse on the rest of the points with a different set of colors. It is easy to show that this gives an $O(n^{6/13})$ CF-coloring if such a monotone sequence always exists (see [HS05] for more details).

Let \mathcal{A} be an $O(n^{1/2})$ conflict-free coloring algorithm [HS05]. Our algorithm can be described as follows. Let $r = n^{\frac{5}{13}}$. Grid the pointset P using G_r such that each row and column has $b = n^{\frac{8}{13}}$ points of P . Compute the dominating sets $D_l(P)$, $1 \leq l \leq 4$ and let $D = \cup_{l=1}^4 D_l(P)$ and $P' = P \setminus D$. We quasi-CF color P' with $\tilde{O}(b^{3/4})$ colors using the algorithm of [EM06] (which uses \mathcal{A} as subroutine). Then, we CF-color D using \mathcal{A} with a different set of colors.

Lemma 3.1 *The above coloring of P is conflict-free.*

Proof. Let $T \in \mathcal{R}$ be a rectangle such that $T \cap P \neq \emptyset$. We show that T contains a point of unique color among the points in $T \cap P$.

We consider 4 cases:

Case 1. A monotone sequence I of size $\Omega(n^{7/13})$ is found: if $T \cap P \setminus I \neq \emptyset$, then by induction and the fact that I and $P \setminus I$ are colored with distinct sets of colors, we know that $T \cap P$ contains a point of a unique color. If $T \cap P \subseteq I$, then $|T \cap P| = 1$ (since I consists of every other point in a monotone sequence) and the statement trivially holds.

We assume thus that case 1 does not hold.

Case 2. $T \cap D \neq \emptyset$: The CF-coloring of D guarantees that there is a point p of unique color among points in $T \cap D$. Since D and $P' = P \setminus D$ are colored with distinct sets of colors, p is a point of unique color among points in $T \cap P$ also.

Case 3. T spans at least 2 rows and 2 columns of G_r : Let (i, j) be a non-empty grid cell of G_r that is intersected by T . Since T contains at least one corner of grid cell (i, j) , $T \cap D_l(R_i \cap C_j) \neq \emptyset$ for some $l \in [1, 4]$, i.e., T contains at least one point of the dominating set of points in grid cell (i, j) . This implies that $T \cap D \neq \emptyset$ and we are back to Case 2.

We may assume now that cases 1, 2 and 3 do not hold.

Case 4. T lies completely within one row or one column of G_r : Since $T \cap P \neq \emptyset$ and $T \cap D = \emptyset$, we have $T \cap P' \neq \emptyset$. The quasi-CF coloring of P' guarantees that there is a point p of unique color among the points in $T \cap P'$. p is also the point of unique color among points in $T \cap P$. \square

Analysis. We bound the total number of colors used by our algorithm. Quasi-CF-coloring of P' requires $\tilde{O}(n^{\frac{8}{13} \times \frac{3}{4}}) = \tilde{O}(n^{6/13})$ colors. To bound the number of colors used in CF-coloring D , we first bound the size of D : $|\mathcal{D}_l^k| = O(n^{7/13})$ for all k , because \mathcal{D}_l^k is a monotonic sequence. Since $D = \cup_{l,h} \mathcal{D}_l^h$ over $1 \leq h \leq 2n^{5/13} - 1$, and $1 \leq l \leq 4$, we have $|D| = O(n^{12/13})$. Thus, the CF-coloring of D (using the $O(n^{1/2})$ -coloring algorithm \mathcal{A}) requires $O(n^{6/13})$ colors. The total number of colors used by our algorithm is thus $\tilde{O}(n^{6/13})$.

Theorem 3.1 *Any pointset $P \subseteq R^2$, can be CF-colored with $\tilde{O}(n^{6/13})$ colors.*

4 Generalized quasi-conflict free coloring

In this section we describe the quasi-CF coloring algorithm. Given an r -grid $G_r(P)$ on pointset P , we start by coloring the points of each column, using a CF-coloring algorithm \mathcal{A} as a black-box. We use the same set of colors for all columns. Then we randomly, and independently for each column, we redistribute the colors on the different color classes of the column. Finally, a recoloring step is applied on each monochromatic set of points in each row, again using algorithm \mathcal{A} as the CF-coloring procedure. When we do the recoloring, we color all sets assigned a color ℓ in the first step using the same set of colors S_ℓ . A formal description of this procedure is given in Figure 1.

Procedure QCFC(P, \mathcal{A}, G_r, S):

Input: A pointset $P \subseteq \mathbb{R}^2$, an $f(\cdot)$ -conflict-free coloring algorithm \mathcal{A}
an r -grid G_r on P , and a set of possible colors $S \subseteq \mathbb{N}^*$

Output: A quasi-conflict-free coloring $\chi : P \mapsto S$ of P with respect to G_r .

1. Let N be a subset of \mathbb{N} of size $B(G_r)$
2. **for** $i = 1, \dots, r$ **do**
3. $\chi_i \leftarrow \mathcal{A}(C_i, N)$
4. Let $\pi : N \mapsto N$ be a random permutation
5. **foreach** $p \in C_i$ **do**
6. $\chi'_i(p) \leftarrow \pi(\chi_i(p))$
7. $\chi' \leftarrow \sum_{i=1}^r \chi'_i$; $h \leftarrow \text{range}(\chi')$
8. Let S_1, \dots, S_h be disjoint subsets of \mathbb{N} of size $B(G_r)$
9. **for** $i = 1, \dots, r$ **do**
10. **for** $\ell = 1, \dots, h$ **do**
11. $P_i^\ell \leftarrow \{p \in R_i : \chi'(p) = \ell\}$
12. $\chi''_{i,\ell} \leftarrow \mathcal{A}(P_i^\ell, S_\ell)$
13. $\chi'' \leftarrow \sum_{i=1}^r \sum_{\ell=1}^h \chi''_{i,\ell}$
14. **return** $\chi' \times \chi''$ (mapped to S)

Figure 1: Quasi-conflict-free coloring of a grid

The following is a straightforward generalization of Theorem 3 in [EM06]. We include the proof in the Appendix A for completeness.

Theorem 4.1 *Given any pointset $P \subseteq \mathbb{R}^2$, a grid $G_r = G_r(P)$ with $B(G_r) = B$ on P , and a conflict-free coloring algorithm \mathcal{A} with guarantee $f(\cdot)$ such that $B \geq 4$ and*

$$r \cdot f(B) \log(\delta B) (\log B)^{(-\log B)/2} \leq \frac{1}{2}, \quad (1)$$

procedure QCFC returns a quasi-conflict-free coloring of $G_r(P)$ using

$$q(B) = f(B) f\left(\frac{2\delta B \log B \log(\delta B)}{f(B)}\right) \quad (2)$$

colors, in expected polynomial time, where δ is the constant given in Remark 5.1.

5 Generalized algorithm

In this section, we generalize the algorithm described in Section 3. Recall that, in our coloring algorithm, we used an $O(n^{1/2})$ "black-box" \mathcal{A} for CF-coloring the dominating set D and the quasi-CF-coloring of P' . As a result we obtained an $\tilde{O}(n^{6/13})$ CF-coloring algorithm. We can improve this coloring further by using now this $\tilde{O}(n^{6/13})$

as a new black-box for CF-coloring the dominating set D and quasi-CF-coloring of P' . An easy calculation shows that the number of colors used is asymptotically smaller than $O(n^{6/13})$.

We can now take this approach (almost) to the limit. This results in a succession of strictly improved sequence of algorithms, $\mathcal{A} = \mathcal{A}_0, \mathcal{A}_1, \mathcal{A}_2, \dots$. For $k = 1, 2, \dots$, the structure of \mathcal{A}_k is similar to the algorithm described in Section 3: Grid the pointset P using G_r , where $r = n^{1-\alpha_k}$, for some α_k . Partition P into dominating set D and $P' = P \setminus D$ and use algorithm \mathcal{A}_{k-1} for CF-coloring D and quasi-CF-coloring P' . We choose the parameter α_k such that both the CF-coloring of D and quasi-CF-coloring of P' balance-out into using an $\tilde{O}(n^{\beta_k})$ colors, for some β_k as small as possible.

To be more precise and describe our coloring procedure formally, we need a few more definitions. Given a coloring $\chi : P \mapsto \mathbb{N}^*$, we denote by $\text{range}(\chi) = \{\chi(p) : p \in P\}$, the set of distinct colors used to color P . Let $f : \mathbb{R}_+ \mapsto \mathbb{R}_+$ be a *monotone sub-linear* function on the positive reals. An $f(\cdot)$ -*conflict-free coloring algorithm* \mathcal{A} takes as an input a pointset $P \subseteq \mathbb{R}^2$, and a set of colors $N \subseteq \mathbb{N}^*$ such that $|N| \geq f(|P|)$, and returns a conflict-free coloring $\chi : P \mapsto N$ of P such that $\text{range}(\chi) \leq f(|P|)$.

Remark 5.1 *It will simplify the analysis to assume, without loss of generality, that there exists a constant $\delta > 0$ such that the size of each color class (that is $\max_\ell |\{p \in P : \chi(p) = \ell\}|$) in the coloring returned by \mathcal{A} is at most $\delta|P|/f(|P|)$. (This can be justified as follows. Let $P' \subseteq P$ be the largest monochromatic set returned by \mathcal{A} when applied to P . If $|P'| \geq \delta n/f(n)$, where $n = |P|$, then let P'' be a subset of P' of size exactly $\delta n/f(n)$. Color all points of P'' with the same color ℓ , then the points $P - P''$ recursively with colors different from ℓ . Since $f(n)$ is sublinear, we get the required bound.)*

For disjoint subsets $P', P'' \subseteq P$ and colorings $\chi' : P' \mapsto \mathbb{N}^*$ and $\chi'' : P'' \mapsto \mathbb{N}^*$, we let $\chi' + \chi''$ denote the coloring $\chi : P' \cup P'' \mapsto \mathbb{N}^*$ defined by $\chi(p) = \chi'(p)$ if $p \in P'$ and $\chi(p) = \chi''(p)$ if $p \in P''$. For two colorings $\chi', \chi'' : P \mapsto \mathbb{N}^*$, we denote by $\chi' \times \chi''$ the coloring $\chi : P \mapsto \mathbb{N}^*$ given by $\chi(p) = (\chi'(p), \chi''(p))$ for $p \in P$.

The generalized coloring algorithm is given in Figure 2. Let $f_k(n)$ be the number of colors required by our algorithm $\mathcal{A}_k(P, S)$. We set the values of α_k, β_k, n_k for $k \geq 1$, by the following recurrence relations and formulas:

$$\beta_0 = 1/2, \quad \beta_k = \frac{2\beta_{k-1}(2 - \beta_{k-1})}{3 + \beta_{k-1} - \beta_{k-1}^2}, \quad \beta = (3 - \sqrt{5})/2 \quad (3)$$

$$\alpha_k = \frac{2 - \beta_{k-1}}{3 - \beta_{k-1}} \quad (4)$$

$$n_k = C_0^{\frac{2^k - 1}{\alpha_k}}, \quad C_0 = 5 \quad (5)$$

The structure of the generalised coloring algorithm is the same as the algorithm described in Section 3. Hence, by Lemma 3.1, the above coloring is conflict-free.

Procedure $\mathcal{A}_k(P, S)$:

Input: A pointset $P \subseteq \mathbb{R}^2$, $|P| = n$, a set of colors $S \subseteq \mathbb{N}^*$ of size $f_k(n)$

Output: A conflict-free coloring $\chi : P \mapsto S$ with $|\text{range}(\chi)| \leq f_k(n)$

1. **if** $k = 0$ or $n \leq n_k$ **then**
2. Color P using the \sqrt{n} coloring algorithm
3. **else**
4. Compute α_k and β_k using (3)-(4); Set $r \leftarrow n^{1-\alpha_k}$
5. **if** there is a horizontal or vertical line L containing more than $n_k^{\beta_k} n^{1-\beta_k}$ points of P **then**
6. Let I be the set consisting of every other point of $P \cap L$
7. Color every point of I with the same color $i \in S$
8. $\mathcal{A}_k(P \setminus I, S \setminus \{i\})$
9. **else**
10. Grid P using G_r
11. Compute the dominating set union D w.r.t. $G_r(P)$
12. **if** the longest monotonic sequence L along a diagonal in D is of size $n_k^{\beta_k} n^{1-\beta_k}$ **then**
13. Let I be the set consisting of every other point of L
14. Color every point of I with the same color $i \in S$
15. $\mathcal{A}_k(P \setminus I, S \setminus \{i\})$
16. **else**
17. $\chi' \leftarrow \text{QCFC}(P \setminus D, G_r, \mathcal{A}_{k-1}, S')$
18. $\chi'' \leftarrow \mathcal{A}_{k-1}(D, S \setminus \text{range}(\chi'))$
19. **return** $\chi' + \chi''$

Figure 2: Conflict-free coloring

6 Analysis

Let $f_k(\cdot)$ denote the upper bound on the number of colors required by the algorithm at the k^{th} level. If $n \leq n_k$ or $k = 0$, we use a \sqrt{n} coloring algorithm. Thus, $f_k(n) \geq \sqrt{n}$ for $n \leq n_k$ or $k = 0$. Otherwise, if any one of the dominating sets is larger than $2n_k^{\beta_k} n^{1-\beta_k}$, we color every alternate node in the monotonically increasing dominating set with a single color and recurse on the rest. Thus, $f_k(n) \geq 1 + f_k(n - n_k^{\beta_k} n^{1-\beta_k})$. If there is no such dominating set, we grid the point sets such that no row or column contains more than n^{α_k} points, recursively color the dominating set D and quasi color $P \setminus D$. As stated in Theorem 4.1, if $r \cdot f_{k-1}(K) \log \delta K (\log K)^{-\log K} \leq 1/2$, then quasi coloring a grid of r rows and r columns such that each row and each column contains at most K points requires $f_{k-1}(K) \cdot f_{k-1}(\frac{2\delta K \log K \log \delta K}{f_{k-1}(K)})$ colors. This leads to the following recursion:

$$f_k(n) \geq 4f_{k-1}(n^{(2-\alpha_k-\beta_k)}) + f_{k-1}(n^{\alpha_k}) \cdot f_{k-1}\left(\frac{2\delta n^{\alpha_k} \log n^{\alpha_k} \log \delta n^{\alpha_k}}{f_{k-1}(n)}\right) \quad (6)$$

Let, $\gamma_k = 2^{k+1}$, and $C_k = C_0^{2^k - 1}$, $k > 1$.

Our main theorem in this Section is that $f_k(n) = C_k n^{\beta_k} \log^{\gamma_k} n$ satisfies all the recursions mentioned above. Coupled with Lemma 6.1 where we prove that for $k = O(\log 1/\epsilon)$, $\beta < \beta_k \leq \beta + \epsilon$, this shows that the number of colors required by our algorithm is bounded above by $2^{O(1/\epsilon)} n^{\beta+\epsilon} \log^{O(1/\epsilon)} n = \tilde{O}(n^{\beta+\epsilon})$ for any $\epsilon > 0$.

Lemma 6.1 $\beta < \beta_k \leq \beta + \epsilon$ for $k = O(\log 1/\epsilon)$

Proof. Let us define $\epsilon_k = \beta_k - \beta$. We show that $\epsilon_k = \epsilon$ for $k = \log \frac{2-\beta}{2-2\beta} \frac{0.5-\beta}{\epsilon}$. By definition,

$$\begin{aligned} \beta_k &= \frac{2(\beta + \epsilon_{k-1})(2 - \beta - \epsilon_{k-1})}{3 + (\beta + \epsilon_{k-1}) - (\beta + \epsilon_{k-1})^2} \\ &= \frac{2\beta(2 - \beta) + 2\epsilon_{k-1}(1 - \beta) - \epsilon_{k-1}^2}{(3 + \beta - \beta^2) + (\epsilon_{k-1} - 2\beta\epsilon_{k-1} - \epsilon_{k-1}^2)} \\ &\leq \frac{2\beta(2 - \beta) + 2\epsilon_{k-1}(1 - \beta)}{(3 + \beta - \beta^2) + (\epsilon_{k-1} - 2\beta\epsilon_{k-1})} \\ &\leq \frac{2\beta(2 - \beta) + 2\epsilon_{k-1}(1 - \beta)}{(3 + \beta - \beta^2)} \quad (\text{assuming } 2\beta \leq 1) \end{aligned}$$

Since $3 + \beta - \beta^2 = 2(2 - \beta)$, it implies

$$\beta_k \leq \beta + \frac{4\epsilon_{k-1}(1 - \beta)}{2(2 - \beta)}$$

Therefore,

$$\epsilon_k \leq \frac{\epsilon_{k-1}(2 - 2\beta)}{(2 - \beta)}$$

Since this is true for any $k > 0$, we get

$$\begin{aligned} \epsilon_k &\leq \left(\frac{2 - 2\beta}{2 - \beta}\right)^k \epsilon_0 \\ &= \left(\frac{2 - 2\beta}{2 - \beta}\right)^k (0.5 - \beta) \end{aligned}$$

Thus for $k = \log \frac{2-\beta}{2-2\beta} \frac{0.5-\beta}{\epsilon}$, $\epsilon_k \leq \epsilon$.

Lemma 6.2 *There exists constants (w.r.t. n , but dependent on k) C_k, γ_k, n_k such that they satisfy the following recurrences:*

$$C_k \geq 4C_{k-1} + C_{k-1}^{2-\beta_{k-1}} \quad (\text{for all } k > 0) \quad (7)$$

$$C_k \geq ((n_k^\beta - (n - n_k^{\beta_k} n^{1-\beta_k})^{\beta_k}) \log^{\gamma_k} n)^{-1} \quad (\text{for all } k > 0 \text{ and } n \geq n_k) \quad (8)$$

$$C_k \geq n_k^{0.5-\beta_k} \log n_k^{-\gamma_k} \quad (\text{for all } k) \quad (9)$$

$$C_k \leq \frac{n_k^{\alpha_k(1+\log \log n_k^{\alpha_k} - \beta_{k-1}) - 1}}{2 \log^{\gamma_k}(n_k^{\alpha_k}) \log(n_k^{\alpha_k}/\beta)} \quad (\text{for all } k) \quad (10)$$

$$\gamma_k \geq \gamma_{k-1}(2 - \beta_{k-1}) + 2\beta_{k-1} \quad (\text{for all } k > 0) \quad (11)$$

It can be easily verified that the constants $C_k = 5^{2^k - 1}$, $\gamma_k = 2^{k+1}$, $n_k = 5^{(2^k - 1)/\alpha_k}$ satisfies the above relations.

Theorem 6.1 $f_k(n) = C_k n^{\beta_k} \log^{\gamma_k} n$ satisfies all of the following recursions:

$$f_k(n) \geq \sqrt{n} \quad (\text{for all } k > 0, n \leq n_k \text{ or } k = 0) \quad (12)$$

$$f_k(n) \geq 1 + f_k(n - n_k^{\beta_k} n^{1-\beta_k}) \quad (\text{for } k > 0 \text{ and } n > n_k) \quad (13)$$

$$n^{1-\alpha_k} f_k(n^{\alpha_k}) \log(n^{\alpha_k}/\beta) (\log n^{\alpha_k})^{-\log n^{\alpha_k}} \leq 1/2 \quad (\text{for } k > 0 \text{ and } n > n_k) \quad (14)$$

$$f_k(n) \geq 4f_{k-1}(n^{2-\alpha_k-\beta_k}) + f_{k-1}(n^{\alpha_k}) \cdot f_{k-1}\left(\frac{2\delta n^{\alpha_k} \log n^{\alpha_k} \log \delta n^{\alpha_k}}{f_{k-1}(n)}\right) \quad (\text{for } k > 0 \text{ and } n > n_k) \quad (15)$$

Proof. Equation 9 states that for all k , $C_k \geq n_k^{0.5-\beta_k} \log n_k^{-\gamma_k}$. This implies that $C_k n_k^{\beta_k} \log n_k^{\gamma_k} \geq \sqrt{n_k}$ and since $C_k n^{\beta_k} \log n^{\gamma_k} - \sqrt{n}$ is a decreasing function of n , $f_k(n) \geq \sqrt{n}$ is true for all $n \leq n_k$. For $k = 0$, $\beta_k = 1/2$ and therefore, Equation 12 is trivially satisfied.

In order to prove Equation 13, we use induction on n . The equation is trivially true for $n = n_k$. By induction hypothesis, we assume that $f_k(n) = C_k n^{\beta_k} \log^{\gamma_k} n$ is true for all $n < r$ and we prove it for r . For this we need to prove that $C_k r^{\beta_k} \log^{\gamma_k} r \geq 1 + C_k (r - n_k^{\beta_k} r^{1-\beta_k})^{\beta_k} \log^{\gamma_k} r$. Equation 8 states that for all $k > 0$ and $r \geq n_k$, $C_k \geq ((r^{\beta_k} - (r - n_k^{\beta_k} r^{1-\beta_k})^{\beta_k}) \log^{\gamma_k} r)^{-1}$. This implies that $C_k (r^{\beta_k} - (r - n_k^{\beta_k} r^{1-\beta_k})^{\beta_k}) \log^{\gamma_k} r \geq 1$ and therefore, $C_k r^{\beta_k} \log^{\gamma_k} r \geq 1 + C_k (r - n_k^{\beta_k} r^{1-\beta_k})^{\beta_k} \log^{\gamma_k} r$, thereby proving Equation 13.

Lemma 6.3 uses Equation 10 to prove Equation 14. Equation 14 implies that the pre-requisite for applying quasi-coloring is satisfied for all $k > 0$ and $n > n_k$.

Lemma 6.4 uses Equation 7 and 11 to prove Equation 15, thereby completing the proof.

Lemma 6.3 For all $k > 0$ and $n > n_k$, $f_k(n) = C_k n^{\beta_k} \log^{\gamma_k} n$ satisfies $n^{1-\alpha_k} f_k(n^{\alpha_k}) \log(n^{\alpha_k}/\beta) (\log n^{\alpha_k})^{-\log n^{\alpha_k}} \leq 1/2$

Proof. We first prove it for $n = n_k$. Equation 10 states that

$$C_k \leq \frac{n_k^{\alpha_k(1+\log \log n_k^{\alpha_k} - \beta_{k-1}) - 1}}{2 \log^{\gamma_k}(n_k^{\alpha_k}) \log(n_k^{\alpha_k}/\beta)} \quad \text{for all } k. \quad \text{Thus, } \frac{C_k \log^{\gamma_k} \log n_k^{\alpha_k}/\beta}{n_k^{\alpha_k(1+\log \log n_k^{\alpha_k} - \beta_{k-1}) - 1}} \leq 1/2.$$

So, $n_k^{1-\alpha_k} C_k (n_k^{\alpha_k})^{\beta_k} \log^{\gamma_k}(n_k^{\alpha_k}) \log(n_k^{\alpha_k}/\beta) n_k^{-\log \log n_k^{\alpha_k}} \leq 1/2$.

Thus, $n_k^{1-\alpha_k} C_k (n_k^{\alpha_k})^{\beta_k} \log^{\gamma_k}(n_k^{\alpha_k}) \log(n_k^{\alpha_k}/\beta) (\log(n_k^{\alpha_k}))^{-\log(n_k^{\alpha_k})} \leq 1/2$.

Since $n^{1-\alpha_k} C_k (n^{\alpha_k})^{\beta_k} \log^{\gamma_k}(n^{\alpha_k}) \log(n^{\alpha_k}/\beta) (\log(n^{\alpha_k}))^{-\log(n^{\alpha_k})}$ is a decreasing function of n , this is true for all $n > n_k$.

Lemma 6.4 For all $k > 0$ and $n \geq n_k$, $f_k(n) = C_k n^{\beta_k} \log^{\gamma_k} n$ satisfies $f_k(n) \geq 4f_{k-1}(n^{2-\alpha_k-\beta_k}) + f_{k-1}(n^{\alpha_k}) \cdot f_{k-1}\left(\frac{2\delta n^{\alpha_k} \log n^{\alpha_k} \log \delta n^{\alpha_k}}{f_{k-1}(n)}\right)$

The proof of this Lemma can be found in Appendix B

Using Theorem 6.1 and Lemma 6.1, we get the following:

Theorem 6.2 Let P be a set of n points in \mathbb{R}^2 . Algorithm $A_k(P, S)$ conflict-free colors P with respect to rectangle ranges using $2^{O(1/\epsilon)} n^{\beta+\epsilon} \log^{O(1/\epsilon)} n = \tilde{O}(n^{\beta+\epsilon})$ colors, for any arbitrarily small $\epsilon > 0$.

7 Discussion

Note that the quasi-CF-coloring Algorithm QCF when fed with an $\tilde{O}(n^{\beta_{k-1}})$ -CF-coloring algorithm \mathcal{A}_{k-1} as an input, returns a $\tilde{O}(n^{\beta_k}) = \tilde{O}(B(G_r)^{\beta_{k-1}(2-\beta_{k-1})})$ -quasi-CF-coloring of the grid $G_r(P')$, defined on subset $P' \subseteq P$, where $r = n^{1-\alpha_k}$. This is actually how the successive improvements are made, since $\beta_k < \beta_{k-1}$. Suppose there exists a quasi-CF-coloring algorithm that uses $K(G_r)^c$ colors for some $c > 0$. Then an easy calculating shows that our algorithm \mathcal{A}_k returns a CF-coloring using $n^{c/(c+1)}$ colors. Clearly any improvement on the quasi-CF-coloring algorithm will translate to an improvement on the general case. More precisely, we have the following.

Corollary 7.1 Let P be a set of points of size n , and $r = n^{1-\alpha}$ for some $\alpha \in (0, 1)$. If there exists a quasi-CF-coloring algorithm of the grid $G_r(P')$ that requires $O(n^c)$ colors for some $c > 0$ then, we can obtain a CF-coloring algorithm of P that requires $O(n^{c/(c+1)})$ colors.

By setting $c = \epsilon$, we have the following:

Corollary 7.2 Let P be a set of points of size n , and $r = n^{1-\alpha}$ for some $\alpha \in (0, 1)$. If there exists a quasi-CF-coloring algorithm of the grid $G_r(P')$ that requires $O(n^\epsilon)$ colors for any $\epsilon > 0$, then we can obtain a CF-coloring algorithm of P that requires $O(n^\epsilon)$ colors.

One might think that, if such an improved CF-coloring algorithm (that uses $O(n^\beta)$ colors, $\beta < 0.382$) is obtained, it could be further improved using our iterated improvement scheme given in Section 5. This is not possible, since we can easily show that, for any $\beta_{k-1} < 0.382$, there exists no $\alpha_k \in (0, 1)$ for which the quasi-CF coloring of P' and CF-coloring of dominating set D can both be done using $O(n^{\beta_k})$ colors where $\beta_k < \beta_{k-1}$. Thus, one cannot hope to obtain a CF-coloring algorithm that uses fewer than $O(n^{.382})$ colors by using our idea.

A Appendix: Proof of Generalized quasi-conflict free coloring theorem

Theorem A.1 Given any pointset $P \subseteq \mathbb{R}^2$, a grid $G_r = G_r(P)$ with $B(G_R) = B$ on P , and a conflict-free coloring algorithm \mathcal{A} with guarantee $f(\cdot)$ such that $B \geq 4$ and

$$r \cdot f(B) \log(\delta B) (\log B)^{(-\log B)/2} \leq \frac{1}{2}, \quad (16)$$

procedure *QCFC* returns a quasi-conflict-free coloring of $G_r(P)$ using

$$q(B) = f(B) f\left(\frac{2\delta B \log B \log(\delta B)}{f(B)}\right) \quad (17)$$

colors, in expected polynomial time, where δ is the constant given in Remark 5.1.

Proof. Let $\chi_i, \chi', \chi'', h, P_i^\ell$ be as defined in the procedure, and $\chi = \chi' \times \chi''$ be the coloring returned in Step 14. The theorem follows from the following two claims.

Claim A.1 ([EM06]) χ is quasi-conflict-free.

Proof. Let $T \in \mathcal{R}$ be any rectangle that lies completely inside a row or a column of G_r , such that $T \cap P \neq \emptyset$. If T contains only points belonging to a single column C_j of G_r , then the fact algorithm \mathcal{A} returns a conflict-free coloring of C_j and the definition of χ'_j imply that T contains a point $p \in T \cap C_j$ such that $\chi'_j(p) \neq \chi'_j(p')$ for all $p \neq p' \in T \cap P$. Then $\chi'(p)$ and hence $\chi(p)$ is different in the first coordinate from $\chi(p')$ for every $p \neq p' \in T \cap P$. Now assume that T contains only points belonging to a single row i of G_r . Since $T \cap P \neq \emptyset$, there is an $\ell \in [m]$ such that $T \cap P_i^\ell \neq \emptyset$. Since \mathcal{A} returns a conflict-free coloring $\chi''_{i,\ell}$ of P_i^ℓ , there is a point $p \in T \cap P_i^\ell$, such that $\chi''_{i,\ell}(p) \neq \chi''_{i,\ell}(p')$ for all $p \neq p' \in T \cap P_i^\ell$. Thus if $p' \in T \cap R_i$, then either $p' \in P_i^{\ell'}$ for $\ell' \neq \ell$ in which case $\chi'(p') \neq \chi'(p)$, or $p' \in P_i^\ell$ but $\chi''(p') \neq \chi''(p)$. In both cases $\chi(p') \neq \chi(p)$. \square

Claim A.2 With probability at least $1/2$, $|\text{range}(\chi)| \leq q(B)$ given by (2).

Proof. Fix $i \in [r]$ and $\ell \in [h]$. Define $t = \delta B/h$. For $j \in [r]$, let $A_{i,j}^\ell = \{p \in C_j : \chi_j(p) = \ell\}$ and note that $|A_{i,j}^\ell| \leq \delta B/f(B) \leq t$ and $h \leq f(B)$. For $m = 1, 2, \dots, \log t$, let

$$\mathcal{A}_{i,j}^m = \{A_{i,j}^\ell : 2^{m-1} \leq |A_{i,j}^\ell| \leq 2^m, \ell = 1, \dots, h\},$$

and note that

$$\sum_{j=1}^r |\mathcal{A}_{i,j}^m| \leq \frac{B}{2^{m-1}}, \quad (18)$$

since the total number of points in row i of G_r is at most B , and each set in $\mathcal{A}_{i,j}^m$ has at least 2^{m-1} points.

Note that, for any $j \in [r]$, every point $p \in A_{i,j}^\ell$ gets the same color $\chi'(p)$ in Step 6. Thus we can think of the coloring in Step 6 as of permuting randomly the colors to the sets $A_{i,j}^\ell$, $\ell = 1, \dots, h$, and may use $\chi'(A_{i,j}^\ell)$ to denote the color assigned in Step 6 to all points in $A_{i,j}^\ell$. Let $Y_{i,j}^{m,\ell}$ be the indicator random variable that takes value 1 if and only if there exists a set $S \in \mathcal{A}_{i,j}^m$ with $\chi'(S) = \ell$. Let $Y_i^{m,\ell} = \sum_{j=1}^r Y_{i,j}^{m,\ell}$. Then,

$$\mathbb{E}[Y_{i,j}^{m,\ell}] = \Pr[Y_{i,j}^{m,\ell} = 1] = \frac{|\mathcal{A}_{i,j}^m|}{h}$$

$$\mathbb{E}[Y_i^{m,\ell}] = \sum_{j=1}^r \frac{|\mathcal{A}_{i,j}^m|}{h} \leq \frac{B}{h2^{m-1}} = \frac{t}{2^{m-1}},$$

where the last inequality follows from (18).

Note that the variable $Y_i^{m,\ell}$ is the sum of independent Bernoulli trials, and thus applying the Chernoff bound¹, we get

$$\Pr[Y_i^{m,\ell} > \frac{t \log B}{2^{m-1}}] \leq e^{-\frac{t \log B}{4 \cdot 2^{m-1}} \ln\left(\frac{t \log B}{\mathbb{E}[Y_i^{m,\ell}] \cdot 2^{m-1}}\right)}. \quad (19)$$

Using $\mathbb{E}[Y_m^{i,\ell}] \leq t/2^{m-1}$ and $2^m \leq t$, we deduce from (19) that

$$\Pr[Y_i^{m,\ell} > \frac{t \log B}{2^{m-1}}] \leq (\log B)^{-(\log B)/2}.$$

Thus, the probability that there exist i , ℓ , and m such that $Y_i^{m,\ell} > t \log B / 2^{m-1}$ is at most

$$rh(\log t)(\log B)^{-(\log B)/2} \leq \frac{1}{2},$$

by (1). Therefore with probability at least $1/2$, $Y_i^{m,\ell} \leq t \log B / (c \cdot 2^{m-1})$ for all i , ℓ , and m . In particular, with constant probability, for all i and ℓ , we have

$$|P_i^\ell| \leq \sum_{m=1}^{\log t} Y_i^{m,\ell} \cdot 2^m \leq 2t \log B \log t.$$

Since algorithm \mathcal{A} has guarantee $f(\cdot)$, with constant probability, the total number of colors needed, by the sublinearity of $f(\cdot)$, is

$$|\text{range}(\chi)| \leq \sum_{\ell=1}^h f(|P_i^\ell|) \leq h \cdot f(2t \log B \log t) \leq q(B),$$

as claimed. \square

¹In particular, the following version [MR95]: $\Pr[X \geq (1 + \delta)\mu] \leq e^{-(1+\delta) \ln(1+\delta)\mu/4}$, for $\delta > 1$ and $\mu = \mathbb{E}[X]$.

B Appendix: Proof of Lemma 6.4

Lemma B.1 For all $k > 0$ and $n \geq n_k$, $f_k(n) = C_k n^{\beta_k} \log^{\gamma_k} n$ satisfies $f_k(n) \geq 4f_{k-1}(n^{2-\alpha_k-\beta_k}) + f_{k-1}(n^{\alpha_k}) \cdot f_{k-1}(\frac{2\delta n^{\alpha_k} \log n^{\alpha_k} \log \delta n^{\alpha_k}}{f_{k-1}(n)})$

Proof. We prove by induction on k . For $k = 0$ the claim is true since we always have a $\sqrt{n} \log n$ coloring. Assuming it to be true for all $k < k'$, we need to prove that $C_{k'} n^{\beta_{k'}} \log^{\gamma_{k'}} n \geq g(n)$ where $g(n)$ is given by the following equation:

$$g(n) = 4C_{k'-1}(n^{(2-\alpha_{k'}-\beta_{k'})})^{\beta_{k'-1}} \log^{\gamma_{k'-1}}(n^{(2-\alpha_{k'}-\beta_{k'})}) + C_{k'-1} n^{\alpha_{k'} \beta_{k'-1}} \log^{\gamma_{k'-1}} n^{\alpha_{k'}} \cdot C_{k'-1} \left(\frac{2\delta n^{\alpha_{k'}} \log n^{\alpha_{k'}} \log \delta n^{\alpha_{k'}}}{C_{k'-1} n^{\alpha_{k'} \beta_{k'-1}} \log^{\gamma_{k'-1}} n^{\alpha_{k'}}} \right)^{\beta_{k'-1}} \log^{\gamma_{k'-1}} \left(\frac{2\delta n^{\alpha_{k'}} \log n^{\alpha_{k'}} \log \delta n^{\alpha_{k'}}}{C_{k'-1} n^{\alpha_{k'} \beta_{k'-1}} \log^{\gamma_{k'-1}} n^{\alpha_{k'}}} \right)$$

Since $\delta = 1/\beta$

$$\begin{aligned} g(n) &\leq 4C_{k'-1}(n^{(2-\alpha_{k'}-\beta_{k'})})^{\beta_{k'-1}} \log^{\gamma_{k'-1}} n \\ &\quad + (C_{k'-1} n^{\alpha_{k'} \beta_{k'-1}} \log^{\gamma_{k'-1}} n^{\alpha_{k'}})^{1-\beta_{k'-1}} \cdot C_{k'-1} (n^{\alpha_{k'}} \log^2 n^{\alpha_{k'}})^{\beta_{k'-1}} \log^{\gamma_{k'-1}} n \\ &\leq 4C_{k'-1} n^{(2-\alpha_{k'}-\beta_{k'})\beta_{k'-1}} \log^{\gamma_{k'-1}} n \\ &\quad + (C_{k'-1})^{2-\beta_{k'-1}} n^{\alpha_{k'} \beta_{k'-1} (2-\beta_{k'-1})} \log^{((2-\beta_{k'-1})(\gamma_{k'-1})+2\beta_{k'-1})} n \end{aligned}$$

Using Equation 11 in the above inequality, we get the following:

$$\begin{aligned} g(n) &\leq 4C_{k'-1} n^{(2-\alpha_{k'}-\beta_{k'})\beta_{k'-1}} \log^{\gamma_{k'}} n \\ &\quad + (C_{k'-1})^{2-\beta_{k'-1}} n^{\alpha_{k'} \beta_{k'-1} (2-\beta_{k'-1})} \log^{\gamma_{k'}} n \end{aligned}$$

Since by our definition of $\alpha'_k, \beta'_k, (2-\alpha'_k-\beta'_k)\beta_{k'-1} \leq \beta'_k, \alpha'_k \beta_{k'-1} (2-\beta_{k'-1}) \leq \beta'_k$,

$$g(n) \leq (4C_{k'-1} + C_{k'-1}^2) n^{\beta_{k'}} \log^{\gamma_{k'}} n$$

Using Equation 7 in the above inequality, we get $g(n) \leq C_{k'} n^{\beta_{k'}} \log^{\gamma_{k'}} n$, thus completing the proof.

References

- [AKS99] N. Alon, M. Krivelevich, and B. Sudakov. Coloring graphs with sparse neighborhoods. *J. Combinatorial Theory Ser. B.*, 77:73–82, 1999.
- [AS06] Noga Alon and Shakhar Smorodinsky. Conflict-free colorings of shallow discs. In *SCG '06: Proceedings of the twenty-second annual symposium on Computational geometry*, pages 41–43, New York, NY, USA, 2006. ACM Press.

- [BNCS06] Amotz Bar-Noy, Panagiotis Cheilaris, and Shakhar Smorodinsky. Conflict-free coloring for intervals: from offline to online. In *SPAA '06: Proceedings of the eighteenth annual ACM symposium on Parallelism in algorithms and architectures*, pages 128–137, New York, NY, USA, 2006. ACM Press.
- [Che06] Ke Chen. How to play a coloring game against a color-blind adversary. In *SCG '06: Proceedings of the twenty-second annual symposium on Computational geometry*, pages 44–51, New York, NY, USA, 2006. ACM Press.
- [CKS06] Ke Chen, Haim Kaplan, and Micha Sharir. Online cf coloring for half-planes, congruent disks, and axis-parallel rectangles. *Manuscript*, 2006.
- [ELRS03] G. Even, Z. Lotker, D. Ron, and S. Smorodinsky. Conflict-free colorings of simple geometric regions with applications to frequency assignment in cellular networks. *SIAM J. Comput.*, 33:94–136, 2003.
- [EM06] Khaled M. Elbassioni and Nabil H. Mustafa. Conflict-free colorings of rectangles ranges. In *STACS*, pages 254–263, 2006.
- [FLM⁺05] A. Fiat, M. Levy, J. Matousek, E. Mossel, J. Pach, M. Sharir, S. Smorodinsky, U. Wagner, and E. Welzl. Online conflict-free coloring for intervals. In *Proc. 16th Annual ACM-SIAM Symposium on Discrete Algorithms*, 2005.
- [HS05] S. Har-Peled and S. Smorodinsky. Conflict-free coloring of points and simple regions in the plane. *Discrete & Comput. Geom.*, 34:47–70, 2005.
- [MR95] R. Motwani and P. Raghavan. *Randomized Algorithms*. Cambridge University Press, New York, NY, 1995.
- [PT03] J. Pach and G. Toth. Conflict free colorings. In *Discrete & Comput. Geom., The Goodman-Pollack Festschrift*. Springer Verlag, Heidelberg, 2003.
- [Smo06] Shakhar Smorodinsky. On the chromatic number of some geometric hypergraphs. In *SODA '06: Proceedings of the seventeenth annual ACM-SIAM symposium on Discrete algorithm*, pages 316–323, New York, NY, USA, 2006. ACM Press.